



Docket No. 01211.001US1
WD # 369724.wpd

SALK No. S99046/PC

Clean Version of Pending Claims

**SEQUENCE AND METHOD FOR INCREASING PROTEIN EXPRESSION IN CELLULAR
EXPRESSION SYSTEMS**

Applicant: Martin Latterich et al.

Serial No.: 09/458,779

-
1. (Cancel) A polynucleotide comprising a sequence encoding a functional vesicular fusion factor 2 protein (Vff2p), or a structural homolog of Vff2p.
 2. (Cancel) The polynucleotide of claim [1] 3, wherein the Vff2p is a yeast protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion.
 3. (Amended) A polynucleotide comprising a sequence encoding a functional vesicular fusion factor 2 protein (Vff2p), wherein the Vff2p comprises SEQ ID NO:2 or a variant thereof, and wherein the Vff2p is a yeast protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion.
 4. (Amended) The polynucleotide of claim 3, comprising SEQ ID NO:1 or a variant thereof.
 5. (Amended) The polynucleotide of claim 3, wherein the protein is about 32 kD.
 6. (Amended) The polynucleotide of claim 3, further comprising a promoter operatively linked to the sequence encoding the Vff2p.
 7. The polynucleotide of claim 6 wherein the promoter is a promoter that functions in a host cell to direct transcription of the sequence encoding the Vff2p.
 8. (Amended) The polynucleotide of claim 3, further comprising a sequence encoding a heterologous target protein.

9. The polynucleotide of claim 8, further comprising a second promoter operably linked to the sequence encoding the target protein.
10. The polynucleotide of claim 9, wherein the second promoter is a promoter that functions in the host cell to direct transcription of the target protein.
11. The polynucleotide of claim 7, wherein the host cell is a yeast cell.
12. (Amended) The polynucleotide of claim 2, wherein the yeast is a *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Yarrowia lipolytica*, *Pichia pastoris*, *Hansenula polymorpha*, or *Kluyveromyces lactis*.
13. [Cancel] A polynucleotide expression vector comprising a polynucleotide encoding a functional Vff2p, or a structural homolog of Vff2p.
14. (Amended) A polynucleotide expression vector comprising a polynucleotide encoding a functional Vff2p, wherein the Vff2p comprises SEQ ID NO:2, or a variant thereof, and wherein the Vff2p is a yeast protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion.
15. (Amended) The expression vector of claim 14, comprising SEQ ID NO:1, or a variant thereof.
16. (Amended) The expression vector of claim 14, wherein the protein is about 32 kD.

17. (Amended) The expression vector of claim 14, further comprising a promoter sequence operatively linked to the sequence encoding the Vff2p.
18. The expression vector of claim 17 wherein the promoter is a promoter that functions in a host cell to direct transcription of the sequence encoding the Vff2p.
19. (Amended) The expression vector of claim 14, further comprising a sequence encoding a heterologous target protein.
20. The expression vector of claim 19, wherein transcription of the target protein is directed by a second promoter.
21. The expression vector of claim 20, wherein the second promoter is a promoter that functions in the host cell to direct transcription of the target protein.
22. The expression vector of claim 18, wherein the host cell is a yeast cell.
23. The expression vector of claim 22, wherein the yeast is *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Yarrowia lipolytica*, *Pichia pastoris*, *Hansenula polymorpha*, or *Kluyveromyces lactis*.
24. (Cancel) A recombinant yeast cell comprising a cell genetically altered to express a protein encoded by a polynucleotide sequence encoding a functional Vff2p wherein the Vff2p increases cell mass and secretion of expressed proteins in yeast.

25. (Amended) A recombinant host cell comprising a yeast cell genetically altered to express a protein encoded by a polynucleotide sequence encoding a functional Vff2p, wherein the Vff2p comprises SEQ ID NO:2, or a variant thereof, and wherein the Vff2p is a yeast protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion.
26. (Amended) The host cell of claim 25, comprising SEQ ID NO:1, or a variant thereof.
27. The host cell of claim 25, further comprising a sequence encoding a heterologous target protein.
28. (Cancel) The host cell of claim 24, wherein the host cell is a yeast cell.
29. (Amended) The host cell of claim 25, wherein the yeast cell is a *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Yarrowia lipolytica*, *Pichia pastoris*, *Hansenula polymorpha*, or *Kluyveromyces lactis* cell.
30. (Amended) The host cell of claim 25, wherein the host cell lacks a functional protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion, other than Vff2p.
31. (Amended) A method for increasing cell growth of a host cell, comprising introducing Vff2p into the cell and culturing the cell, wherein the Vff2p has at least 40% homology to SEQ ID NO:2.

32. (Amended) The method for increasing cell growth of a cell according to claim 31, wherein the host cell is cultured under conditions effective to allow expression of the encoded Vff2p.
33. (Amended) A method for increasing protein secretion from a host cell, comprising introducing Vff2p into the cell and culturing the cell, wherein the Vff2p has at least 40% homology to SEQ ID NO:2.
34. (Amended) The method for increasing protein secretion from a cell according to claim 33, wherein the host cell is cultured under conditions effective to allow expression of the encoded Vff2p.
35. (Cancel) An isolated functional vesicular fusion factor 2 protein.
36. (Amended) An isolated functional vesicular fusion factor 2 protein comprising SEQ ID NO:2, or a variant thereof, and wherein the Vff2p is a yeast protein involved in the secretory pathway and/or involved in the required cellular machinery for membrane fusion.
37. (Amended) A method of selecting for a yeast secretory mutant cell containing a polynucleotide sequence encoding a Vff2p operably linked to a promoter, wherein the Vff2p comprises SEQ ID NO:2, or a variant thereof, the method comprising growing the secretory mutant cell at a restrictive temperature of about 32-37°C, wherein the restrictive temperature selectively favors mutant cell growth.
38. The method of claim 37, wherein the temperature is at about 37°C.

39. The method of claim 37, wherein the secretory mutant cell is sec17-1, sec18-1, bet1-1, sec22-2, usol-1, pex3-1, sed5-1, cdc48-2, sec7-5, or ypt1-3.28.
40. The method of claim 39, wherein the secretory mutant cell is sec17-1, sec18-1, bet1-1, sec22-2, usol-1, or pex3-1.
41. The method of claim 40, wherein the secretory mutant cell is sec18-1.
42. The method of claim 37, wherein the polynucleotide further comprises a sequence encoding a heterologous target protein operably linked to a second promoter.
43. The polynucleotide of claim 3, wherein the Vff2p comprises SEQ ID NO:2.
44. (New) The protein of claim 25, wherein the yeast is *S. cerevisiae*.
45. (New) The protein of claim 36, wherein the protein is from *S. cerevisiae*.
46. (New) The method of claim 37, wherein the yeast cell is a *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Yarrowia lipolytica*, *Pichia pastoris*, *Hansenula polymorpha*, or *Kluyveromyces lactis* cell.